



Wyoming

Fazzi Associates Quality Improvement Chart
OASIS-C Home Health Process Measures

Data Collection Period January 2010 to June 2010

Measure	Your Agency	Your State	National Average	National Top 20%
Patient was assessed for pain, using a standardized pain assessment tool.		93%	96%	100%
Patient had pain and pain interventions were included during the care plan and implemented.		92%	94%	100%
Patients exhibited symptoms of heart failure for whom appropriate actions were taken.		95%	97%	100%
Patient was assessed for risk of developing pressure ulcers.		93%	95%	100%
Interventions to prevent pressure ulcers were included in the physician-ordered plan of care for patients assessed to be at risk.		86%	88%	98%
Interventions to prevent pressure ulcers were included in the physician-ordered plan of care and implemented.		85%	86%	98%
The start or resumption of care date was either on the physician-specified date or within 2 days of the referral date.		85%	86%	96%
Patient/ caregiver was instructed on how to monitor the effectiveness of drug therapy, recognize potential adverse effects, and how and when to report problems.		79%	83%	96%
Patients 65 and older had a multi-factor fall risk assessment.		98%	94%	100%
Patients were screened for depression.		88%	91%	99%
Patients received influenza immunization for the current flu season.		73%	67%	83%
Patients were determined to have ever received Pneumococcal Polysaccharide Vaccine (PPV).		68%	60%	81%
Diabetic foot care and education specified during the physician-ordered care plan was implemented.		76%	83%	96%

800-379-0361
www.fazzi.com

©2010

Source: October 2010 Home Health Compare, www.Medicare.gov